VOL. 553, NOS 1 + 2 JOURNAL OF CHROMATOGRAPHY AUGUST 16, 1991

CONTENTS

18th INTERNATIONAL SYMPOSIUM ON CHROMATOGRAPHY, AMSTERDAM, SEPTEMBER 23–28, 1990, PART II

COLUMN LIQUID CHROMATOGRAPHY

Applications: Pharmacological

High-performance liquid chromatography post-column derivatization with fluorescence detection to study the influence of ambroxol and dipalmitoylphosphatidylcholine levels in rabbit eustachian tube washings
by M. Kitsos, C. Gandini, G. Massolini and E. de Lorenzi (Pavia, Italy) and G. Caccialanza (Pisa, Italy) 1

Analysis of sulthiame in serum by narrow-bore high-performance liquid chromatography. Comparison of direct sample injection with pre-column switching and extrelut extraction
by U. H. Juergens (Bielefeld, Germany) 7

Improved high-performance liquid chromatographic resolution of the geometric isomers of 6-hydroxy-4-(1-hydroxy-1-methylethyl)-1-cyclohexene-1-ethanol and by-products with β-cyclodextrin
by A. Italia, L. Dosi and M. Schiavi (Piacenza, Italy) 15

Clean-up of plasma extracts by gel permeation chromatography during analysis of isosorbide nitrates by capillary gas chromatography
by P. O. Edlund and K. Johansen (Solna, Sweden) 21

High-performance liquid chromatographic method for the determination of RGH-5702 in plasma samples
by P. Pudleiner, M. Kapás and L. Vereczkey (Budapest, Hungary) 29

High-performance liquid chromatographic determination of rifapentine and its metabolite in human plasma by direct injection into a shielded hydrophobic phase column
by E. Riva, R. Merati and L. Cavenaghi (Gerenzano, Italy) 35

Micro-liquid chromatography method for the determination of ciclopiroxolamine after pre-column derivatization in topical formulations
by F. Belliardo (Torino, Italy), A. Bertolino and G. Brandolo (Pianezza, Italy) and C. Lucarelli (Rome, Italy) 41

High-performance liquid chromatographic determination of vinblastine, 4-O-deacetylvinblastine and the potential metabolite 4-O-deacetylvinblastine-3-oic acid in biological fluids
by O. van Tellingen and J. H. Beijnen (Amsterdam, The Netherlands), R. Baurain (Brussels, Belgium) and W. W. ten Bokkel Huinink, H. R. van der Woude and W. J. Nooyen (Amsterdam, The Netherlands) 47

Two-dimensional high-performance liquid chromatography at low ng/ml levels of the anti-proliferative agent B859-35 in serum with automated sample clean-up, solid-phase trapping and ultraviolet detection
by K. Zech and R. Herzog (Konstanz, Germany) 55

Sensitive determination of the phosphodiesterase III/IV inhibitor zardaverine in human serum by direct sample injection, automated precolumn clean-up and high-performance liquid chromatography
by R. Huber and U. Schildnitz (Konstanz, Germany) 65

Supercritical fluid chromatography in the routine stability control of antipruritic preparations
by K. Anton, M. Bach and A. Geiser (Basle, Switzerland) 71

High-performance liquid chromatographic evaluation of Med 15 and its metabolites Med 5 and tolmetin in rat plasma
by A. Mancinelli, G. Bruno, G. Cardace, E. Morabito, A. Marzo and E. Arrigoni Martelli (Rome, Italy) 81

Applications: Proteins and Their Constituents

Standardization of cation-exchange clean-up prior to gas chromatography of amino acids
by V. Fábián, M. Morvai, M. Pintér-Szakács and I. Molnár-Perl (Budapest, Hungary) . . 87

Separation of elastin cross-links as phenylisothiocyanate derivatives
by T. Hanis, Z. Deyl, R. Struzinsky and I. Miksik (Prague, Czechoslovakia) 93

Recent advances in liquid chromatography–mass spectrometry and capillary zone electrophoresis–mass spectrometry for protein analysis
by M. J.-F. Suter, B. B. DaGue, W. T. Moore, S.-N. Lin and R. M. Caprioli (Houston, TX, U.S.A.) 101

Purification of canine prolactin and growth hormone by fast protein liquid chromatography
by C. Giard-Pasquier and U. Rickenbacher (Basle, Switzerland) 117

Advantages and limitations of pre-column derivatization of amino acids with dabsyl chloride
by E. H. J. M. Jansen, R. H. van den Berg, R. Both-Miedema and L. Doorn (Bilthoven, The Netherlands) 123

Comparison of high-performance liquid chromatographic detection methods for thyronine and tyrosine residues in toxicological studies of the thyroid
by L. Doorn, E. H. J. M. Jansen and F. X. R. van Leeuwen (Bilthoven, The Netherlands) 135

Normal- and reversed-phase high-performance liquid chromatography of some phosphonodipeptides
by S. V. Galushko, M. Y. Belik, V. A. Solodenkó, T. N. Kasheva and V. P. Kukhar (Kiev, U.S.S.R.) 143

Rapid and sensitive high-performance liquid chromatographic method for the analysis of tryptophan, tyrosine and phenylalanine in biological samples
by C. Carducci, F. Moretti, M. Birarelli and I. Antonozzi (Rome, Italy) 149

Improvement of extraction and concentration of milk peptides with solid-phase cartridges for analysis by high-performance liquid chromatography
by A. Voirin and J.-F. Letavernier (Le Plessis-Robinson, France) and B. Sebille (Thiais, France) 155

Anion-exchange chromatographic properties of α-lactalbumin eluted from quaternized polyvinylimidazole. Study of the role of the polymer coating
by R. Lemque, C. Vidal-Madjar, M. Racine, J. Piquion and B. Sébille (Thiais, France) . . 165

Other Biological Applications

Determination of several retinoids, carotenoids and E vitamers by high-performance liquid chromatography. Application to plasma and tissues of rats fed a diet rich in either β-carotene or canthaxanthin
by T. van Vliet (Zeist and Amsterdam, The Netherlands) and F. van Schaik, J. van Schoonhoven and J. Schrijver (Zeist, The Netherlands) 179

Application of high-performance liquid chromatography to the analysis of niacin and biotin in Italian almond cultivars
by A. Rizzolo, C. Baldo and A. Polesello (Milan, Italy) 187

Determination of the ultraviolet absorbance and radioactivity of purine compounds separated by high-performance liquid chromatography. Application to metabolite flux rate analysis
by T. Grune, W. G. Siems, G. Gerber and R. Uhlig (Berlin, Germany) 193

Hypertensive congenital adrenal enzymatic defects detected by high-performance liquid chromatography of corticosteroids
by G. Carpenè, A. Vettoretti, F. Pedini and S. Rocco (Padova, Italy), F. Mantero (Catania, Italy) and G. Opocher (Padova, Italy) 201

Purine and pyrimidine compounds in murine peritoneal macrophages cultured *in vitro*
by A. Werner (Berlin, Germany), R. Maeba, H. Shimasaki and N. Ueta (Tokyo, Japan) and G. Gerber (Berlin, Germany) 205

Analysis of biomass pyrolysis oils by a combination of various liquid chromatographic techniques and gas chromatography–mass spectrometry
by P. L. Desbène (Evreux, France), M. Essayegh and B. Desmazieres (Paris, France) and F. Villeneuve (Nogent sur Marne, France) 211

Identification of flavonoids from *Ginkgo biloba* L., *Anthemis nobilis* L. and *Equisetum arvense* L. by high-performance liquid chromatography with diode-array UV detection
by P. Pietta, P. Mauri and A. Bruno (Milan, Italy), A. Rava (Pavia, Italy) and E. Manera and P. Ceva (Mede, Italy) 223

Measurement of enzyme activities of cytochrome P-450 isoenzymes by high-performance liquid chromatographic analysis of products
by E. J. M. Reinerink, L. Doorn, E. H. J. M. Jansen and A. A. J. van Iersel (Bilthoven, The Netherlands) 233

Determination of ethylenediaminetetraacetic acid and its salts in canned mushrooms by reversed-phase ion-pair liquid chromatography
by J. de Jong, A. van Polanen and J. J. M. Driessen (Wageningen, The Netherlands) . . 243

High-performance liquid chromatographic determination of acetone in blood and urine in the clinical diagnostic laboratory
by A. Brega, P. Villa, G. Quadrini and A. Quadri (Brescia, Italy) and C. Lucarelli (Rome, Italy) 249

Selective post-column fluorigenic reaction with benzamidine for trace level detection of reducing saccharides in liquid chromatography
by A. Coquet, J.-L. Veuthey and W. Haerdi (Geneva, Switzerland) 255

Determination of free DOPA and 3-O-methyl-DOPA in human plasma by high-performance liquid chromatography with electrochemical detection
by G. V. Melzi d'Eril and V. Rizzo (Pavia, Italy) 265

High-performance liquid chromatographic determination of sugars and polyols in extracts of lichens and sugarcane juice
by C. Vicente, J. L. Mateos, M. M. Pedrosa and M. Estrella Legaz (Madrid, Spain) . . . 271

Evaluation of different packings for high-performance liquid chromatographic analysis of alkyl lysophospholipids
by J. Coene, M. Ghijs, E. van den Eeckhout, W. van den Bossche and P. Sandra (Ghent, Belgium) 285

Applications: Environmental

Rapid determination of abamectin in lettuce and cucumber by high-performance liquid chromatography
by J. Vuik (Zeist, The Netherlands) 299

Screening for coumatetralyl in soft drinks by solid-matrix extraction and high-performance liquid chromatography with diode-array detection
by A. di Muccio, I. Camoni, L. Vergori, R. Dommarco, D. Attard Barbini, F. Vergori, A. Ausili and A. Santilio (Rome, Italy) 305

Evaluation of eluents in thermospray liquid chromatography–mass spectrometry for identification and determination of pesticides in environmental samples
by D. Barceló and G. Durand (Barcelona, Spain) and R. J. Vreeken, G. J. de Jong, H. Lingeman and U. A. Th. Brinkman (Amsterdam, The Netherlands) 311

Interference removal in the organic trace-level analysis of aqueous environmental samples by on-line liquid chromatographic preconcentration techniques with two precolumns
by V. Coquart and M.-C. Hennion (Paris, France) 329

Sensitive liquid chromatographic determination of alkyl-, nitro- and chlorophenols by precolumn derivatization with dansyl chloride, postcolumn photolysis and peroxyoxalate chemiluminescence detection
by P. J. M. Kwakman, D. A. Kamminga and U. A. Th. Brinkman (Amsterdam, The Netherlands) and G. J. de Jong (Weesp, The Netherlands) 345

Applications: Chiral Separations

Practical strategy for the analytical separation of enantiomers by high-performance liquid chromatography
by R. M. Gaskell and B. Crooks (Macclesfield, U.K.) 357

Influence of chemical structure of tricyclic tertiary dimethylamines on chiral separation by reversed-phase high-performance liquid chromatography after derivatization with (−)-menthylchloroformate
by D. T. Witte, J. Bosman, T. de Boer, B. F. H. Drenth, K. Ensing and R. A. de Zeeuw (Groningen, The Netherlands) 365

(*S,S*)-Diphenylethylethanediamine derivatives as chiral selectors. II. Gasparrini-type bound chiral stationary phase with high enantioselectivity for naphthylamides
by W. Lindner, G. Uray and U. Steiner (Graz, Austria) 373

Reversed-phase and chiral high-performance liquid chromatographic assay of bupivacaine and its enantiomers in clinical samples after continuous extraplural infusion
by B. J. Clark, A. Hamdi, R. G. Berrisford, S. Sabanathan and A. J. Mearns (Bradford, U.K.) 383

Liquid chromatographic separation of the enantiomers of metoprolol and its α-hydroxy metabolite on Chiralcel OD for determination in plasma and urine
by K. Balmér, A. Persson, P.-O. Lagerström and B.-A. Persson (Mölndal, Sweden) and G. Schill (Uppsala, Sweden) 391

Other Applications

Isocratic reversed-phase high-performance liquid chromatographic separation of simple perhalogenated compounds
by P. de A. Leone and C. H. Collins (Campinas, Brazil) 399

Retention of some ethylenediamine oligomers in reversed-phase chromatography
by G. Bánhegyi, T. Cserháti and K. Valkó (Budapest, Hungary) 407

Determination of polymeric hindered amine light stabilizers in polyolefins by high-performance liquid chromatography
by B. Marcato, C. Fantazzini and F. Sevini (Ferrara, Italy) 415

Analysis of polyethylene glycol and derivatives by high-performance liquid chromatography using elevated temperatures and low-wavelength ultraviolet detection, and supercritical fluid chromatography
by R. E. A. Escott (Sunbury-on-Thames, U.K.) and N. Mortimer (Loughborough, U.K.) 423

Micro liquid chromatography with fluorescence detection of thiols and disulphides
by B. Lin Ling (Ghent, Belgium), C. Dewaele (Nazareth, Belgium) and W. R. G. Baeyens (Ghent, Belgium) 433

Quantitative structure-type analysis of heterocompound–hydrocarbon mixtures. VI. Characterization of the nature of neutral heterocompounds in a product of brown coal refining
by H. Zobel, M. Helms and K. Dittberner (Berlin, Germany) 441

PLANAR CHROMATOGRAPHY

Isolation of chlorinated dibenzothiophenes by high-performance thin-layer chromatography
by S. Sinkkonen (Jyväskylä, Finland) 453

Separation of minoxidil and its intermediates by overpressured layer chromatography using a stationary phase bonded with tricaprylmethylammonium chloride
by K. Kovács-Hadady and J. Szilágyi (Debrecen, Hungary) 459

Dependence of the silanophilic effect on the concentration of preadsorbed salts and on the chemical structure of peptides in reversed-phase thin-layer chromatography
by T. Cserháti (Budapest, Hungary) 467

Quantitative two-dimensional thin-layer chromatography
by M. Prošek, I. Drusany and A. Golc-Wondra (Ljubljana, Yugoslavia) 477

Application of high-performance thin-layer chromatography and automated multiple development for the identification and determination of pesticides in water
by U. de la Vigne and D. E. Jänchen (Muttenz, Switzerland) and W. H. Weber (Langenau, Germany) 489

Problems and solutions in chiral thin-layer chromatography: a two-phase "Pirkle" modified amino-bonded plate
by L. Witherow, T. D. Spurway, R. J. Ruane and I. D. Wilson (Macclesfield, U.K.) and K. Longdon (Banbury, U.K.) 497

OTHER SEPARATION METHODS

Chiral separation of cyclodextrin-modified micellar electrokinetic chromatography
by H. Nishi and T. Fukuyama (Osaka, Japan) and S. Terabe (Hyogo, Japan) 503

Non-equilibrium integral Doppler anemometric analysis of particle mixtures in a channel flow using an intrinsic hydrodynamic focusing force biased by another force
by V. L. Kononenko and J. K. Shimkus (Moscow, U.S.S.R.) 517

Author Index Vols. 552 and 553 531

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)